

## Arthropod Phoresy Involving Pseudoscorpions in the Past and Present

George O. Poinar, Jr.<sup>1)</sup>, Božidar P. M. Čurčić<sup>2)</sup>  
and James C. Cokendolpher<sup>3)</sup>

**Abstract** Pseudoscorpions form phoretic associations with a wide range of arthropods, including at least 44 families of insects and three families of arachnids. The present work brings up to date phoretic associations between pseudoscorpions and different arthropods and discusses the long-standing controversy over whether this behavior is basically a predatory response or adapted solely for dispersal. That phoresy in pseudoscorpions is of long standing and obligatory in many cases is demonstrated by its continuance for millions of years, as shown by the fossil record. A member of the Chthoniidae attached to a moth in Baltic amber is reported for the first time.

### Introduction

Pseudoscorpions have the ability to attach themselves to a variety of arthropods, mostly adult stages, which are generally more mobile. The principal benefit of this phoresy is to reach a new habitat with a potentially adequate supply of food. In this way, the distribution of the pseudoscorpions is increased and they are able to exploit transitory habitats that otherwise would be unavailable.

When such phoretic behavior originated is unknown. Although the fossil record of pseudoscorpions extends back to the Middle Devonian, some 380 million years ago (Shear *et al.* 1989), evidence of phoresy between pseudoscorpions and arthropods only appears some 40 million years ago, in the early Tertiary. The present work summarizes knowledge on extant and extinct phoretic relationships between pseudoscorpions and arthropods, discusses the controversy as to whether phoresy is predatory response or an adaptation solely for dispersal, and compares extinct with extant phoretic relationships in regards to the concept of behavioral fixity.

This review recognizes only two of Beier's (1948) three categories of associations between pseudoscorpions and arthropods as providing definite evidence of phoresis. These are when pseudoscorpions are found actually holding the appendages of carriers or when they are riding on the bodies of large arthropods. The third category, pseudoscorpions found in the nests of social insects, while suggestive of phoresy, is considered too vague to be included. However, a new category is introduced here — pseudoscorpions taken from aerial traps on ships and airplanes, indicating phoresy via a winged insect.

---

1) Department of Entomology, Oregon State University, Corvallis, Oregon 97331, USA.

2) Institute of Zoology, Faculty of Biology, University of Belgrade, Studentski Trg 16, 11000 Belgrade, Yugoslavia.

3) 2007 29th Street, Lubbock, Texas 79411, USA.

Accepted July 21, 1998

An attempt is made to cite the original reference to specific examples of phoresy in various geographical areas. Extensive use has been made of previous lists of extant phoretic associations between pseudoscorpions and carriers, especially those of Vachon (1940), Beier (1948), and Muchmore (1971). Multiple references to many of the rider-carrier associations can be found in the above-mentioned works. The present list treats these associations under the carrier arthropod. Terminology follows Beier (1963) and Muchmore (1971).

### Results

Fossil records of arthropod phoresy with pseudoscorpions are limited to examples in Baltic and Dominican ambers (Table 1). Baltic amber is approximately 40 million years old and Dominican amber about 20–40 million years old, although the exact dating of the latter deposits is still controversial (Poinar 1992). A new fossil relationship reported here is a member of the family Chthoniidae attached to the abdominal tergites of a moth in Baltic amber (Fig. 1). The most commonly encountered fossil phoretic associations are between platypodid beetles and pseudoscorpions in Dominican amber (Fig. 2). Extant phoretic associations between pseudoscorpions and arthropods are presented in Table 2.

### Discussion and Conclusions

Direct evidence is provided of four families of pseudoscorpions exhibiting phoretic behavior with eight families of insects in the fossil record and ten families of pseudoscorpions demonstrating phoresy with at least 44 families of insects and three families of arachnids at present. These records will certainly increase as more fossil and recent material is examined.

In the fossil case reported here involving a moth carrying a member of the genus *Chthonius* C. L. Koch (Chthoniidae) in Baltic amber, the pseudoscorpion appears probably to be a juvenile. According to Kaestner (1970) juveniles are never involved in extant phoretic relationships. This agrees with the observations of Zeh and Zeh (1992d), who reported that with *Cordylochernes scorpoides* and the harlequin beetle, "pseudoscorpion presence on beetles is strictly an adult phenomenon". However it appears that juveniles can be phoretic in certain instances, since Sankey (1949) reported two juveniles of *Lamprochernes nodosus* riding on a *Nelima* harvestman, Jones (1978) recorded two nymphs of *Lamprochernes nodosus* attached to *Nelima sylvatica*, and Guilmette *et al.* (1970) reported a pseudoscorpion that was later determined to be a cheliferid protonymph (Harvey 1986) apparently phoretic on a fly in California.

There are two principal methods in which pseudoscorpions can make contact with their carriers in phoretic associations. The most common is when the pseudoscorpion lives in the same habitat as the carrier and makes contact when the latter has completed its development and is ready to leave the old habitat. The less common situation is when the pseudoscorpion waits for a carrier that periodically visits the habitat, such as a bee visiting flowers or collecting resin from a plant (Beier 1948, Muchmore 1971).

Legg (1975) suggested that the reason pseudoscorpions of certain families partake in phoresy more often than those in other families may be associated with the development of a complex mating behavior and the presence of spermathecae for sperm storage.

Table 1. Examples from the fossil record of arthropod phoresy involving pseudoscorpions (D = Dominican amber; B = Baltic amber). Families of pseudoscorpion genera listed: Chthoniidae (*Chthonius*), Cheliferidae (*Dichela*, *Pycnocheilifer*), Chernetidae (*Oligocheernes*), and Withiidae (*Parawithius*).

Carrier	Pseudoscorpion	Amber type	References
INSECTA			
COLEOPTERA			
Platypodidae			
<i>Cenocephalus rhinocerotoides</i>	<i>Parawithius</i> sp.	D	Schlee (1980); Schawaller (1981); Poinar (1992)
DIPTERA			
Rachiceridae			
<i>Chrysothermis speciosa</i>	Unknown	B	Ross (1997)
Tipulidae			
Unknown	Chernetidae	B	Schlee & Glochner (1978)
TRICHOPTERA			
Unknown	<i>Dichela berendtii</i> Menge	B	Beier (1948)
LEPIDOPTERA			
Unknown	<i>Chthonius</i> sp.	B	PRESENT STUDY
HYMENOPTERA			
Apidae			
<i>Proplebeia dominicana</i>	Unknown	D	Wu (1996)
Braconidae			
Unknown	<i>Oligocheernes bachofeni</i> Beier	B	Beier (1937)
Unknown	<i>Pycnocheilifer kleemanni</i> (C. L. Koch et Berendt)	B	Beier (1937)
Ichneumonidae			
Unknown	<i>Dichela berendtii</i> Menge	B	Menge (1855)



Fig. 1. A member of the Chthoniidae (*Chthonius* sp.) on the back of a moth (Lepidoptera) in Baltic amber.

These adaptations have then freed such pseudoscorpions from a purely static existence in permanent habitats and allowed them to visit temporary habitats via phoresy. Legg reasoned that this is why members of the families Chernetidae and Cheliferidae have such a wide distribution.

The carriers can range greatly in size. Some are hardly larger than the pseudoscorpion, while others dwarf the rider. Some large carriers supply not only transportation, but also a survival habitat for the pseudoscorpion; in such cases, the pseudoscorpion may have an extended association with the carrier. An example is the association between pseudoscorpions and the harlequin beetle, *Acrocinus longimanus* (Coleoptera: Cerambycidae). The pseudoscorpions live under the elytra or on other parts of the beetle, often with mites that occur in the same habitat. In the case of *Cordylocheres scorpoides*, one of the six species of pseudoscorpions showing phoretic behavior with this beetle, males use the dorsum of the beetle as a mating site, waiting for new females to climb aboard and occasionally nourishing themselves on some of the associated mites in between periods of mating (Zeh & Zeh 1992d). In the case of these mating males, the acquisition of this habitat may be more for mating than for riding to new sites, and the back of the beetle may be the preferred habitat until that particular male is dislodged by another. That such associations are long-termed is shown by the silken security webs that are produced by the pseudoscorpions on the back of the beetle. The phenomenon of pseudoscorpions spinning silken webs on the back of harlequin beetles was first noted over 80 years ago (Calvert & Calvert 1917) and commented upon more recently (Zeh &




Fig. 2. An adult *Parawithius* sp. (Withiidae) holding a platypodid beetle (*Cenoccephalus rhinocerosoides*) (Platypodidae: Coleoptera) in Dominican amber. The location by which the rider grasps the carrier (under the posterior portion of the elytra) is similar to that reported in extant associations (see illustration in Aguiar *et al.* 1992).

Zeh 1991).

This raises the issue of phoretic pseudoscorpions feeding while riding on their carriers. Consuming mites has been documented with *Cordylochernes scorpiodes* on the harlequin beetle (Zeh & Zeh 1991) and could well occur on larger carriers where the pseudoscorpion assumes a natural position without its chela clasped to an appendages of the carrier. Pseudoscorpions carried by mammals apparently dine on microarthropods, especially mites, in the pelage of their carriers (Muchmore 1971, Durden 1991). It is interesting that some pseudoscorpions found in mammal nests which use the mammal as a carrier are themselves carriers of phoretic mites (Durdin 1991). The presence of phoretic mites on pseudoscorpions carried by arthropods also occurs. One example involved uropodid nymphs on *Titanatelmus conicus*, which in turn was phoretic under the elytra of an *Acanthophorus* sp. (Cerambycidae) in the Belgian Congo (Vachon 1950). Other examples exist, but are not included in the present work.

Vachon (1940) concluded that phoretic behavior in pseudoscorpions was accidental and motivated by a predatory response (hunger for the carrier). Although Muchmore (1971) felt that the reason pseudoscorpions evolved phoretic behavior was unknown, he concluded that it probably was the result of their predaceous nature (attracting them to the carrier). Thus, showing a lack of discrimination, pseudoscorpions would grasp at passing arthropods as prey objects. If the arthropod was large, it would sometimes take

Table 2. Extant associations of arthropod phoresy involving pseudoscorpions. Abbreviations: EU = Europe; AS = Asia; NA = North America; CA = Central America; SA = South America; AU = Australia; NZ = New Zealand; AF = Africa; PA = Pacific Ocean. Families of pseudoscorpion genera listed: Chthoniidae (*Chthonius*, and *Lechytia*), Tridenchthoniidae (*Tridenchthonius*), Neobisiidae (*Neobisium*), Syarinidae (*Syarinus*), Cheiridiidae (*Apocheiridium*, and *Neocheiridium*), Sternophoridae (*Afrostennochorus*), Atemnidae (*Paratennooides*, *Metatenmus*, *Oratenmus*, and *Titanatennus*), Cheliferidae (*Chelifer*, *Ellingsenius*, *Lissochelifer*, *Mucrochelifer*, *Parachelifer*, and *Rhacochelifer*), Chernetidae (*Allochernes*, *Americhernes*, *Anthrenochernes*, *Chelodamus*, *Chernes*, *Cordyllechernes*, *Dinocheirus*, *Hesperochernes*, *Incachernes*, *Lamprochernes*, *Lustrochernes*, *Muscichernes*, *Parachernes*, *Pselaphochernes*, *Semeiochernes*, and *Sphaenochernes*, and *Withius*), *Withiidae* (*Dolichowithius*, *Parawithius*, *Stenowithius*, and *Withius*).

Carrier	Pseudoscorpion	Region	References
<b>ARACHNIDA</b>			
<b>ARANEAE</b>			
Heteropodidae			
<i>Dolena cancerides</i>	<i>Afrostennochorus anabates</i> Harvey	AU	Harvey (1985)
<i>Olios fasciculatus</i>	<i>Lustrochernes grossus</i> (Banks)	NA	Hoff & Jennings (1974)
<b>OPILIONES</b>			
Sclerosomatidae			
<i>Leiobunum blackwalli</i>	<i>Pselaphochernes dubius</i> (O.-P. Cambridge)	EU	Vachon (1947)
<i>Leiobunum rotundum</i>	<i>Pselaphochernes dubius</i> (O.-P. Cambridge)	EU	Vachon (1947)
<i>Nelima gothica</i>	<i>Lamprochernes nodosus</i> (Schränk)	EU	Sankey (1949)
Phalangidae			
<i>Oligolophus meadii</i>	<i>Pselaphochernes dubius</i> (O.-P. Cambridge)	EU	Cloudsley-Thompson (1956)
<i>Opilio parietinus</i>	<i>Lamprochernes nodosus</i> (Schränk)	EU	Sankey (1949)
<i>Mitopus morio</i>	<i>Lamprochernes nodosus</i> (Schränk)	EU	Sankey (1949)
<i>Nelima sylvatica</i>	<i>Lamprochernes nodosus</i> (Schränk)	EU	Jones (1978)
<i>Phalangium opilio</i>	<i>Lamprochernes nodosus</i> (Schränk)	EU	Leydig (1867), Spicer (1867)
<i>Phalangium opilio</i>	<i>Chelifer cancrinoides</i> (Linnaeus)	EU	Spicer (1867)
<b>INSECTA</b>			
<b>ODONATA</b>			
Unknown	Unknown	NA	Dunkle (1984)
<b>ORTHOPTERA</b>			
Acrididae			
Unknown	<i>Lustrochernes concinnus</i> Hoff	SA	Hoff (1947)
Rhaphidophoridae	<i>Chernes cavicola</i> G. Joseph	EU	Joseph (1882)
<i>Troglophilus cavicola</i>			
<b>HEMIPTERA</b>			
Reduviidae	<i>Titanatennus coreophilus</i> Beier	AF	Beier (1948)
<i>Khafra praedo</i>			
<b>COLEOPTERA</b>			
Cerambycidae	<i>Titanatennus congicus</i> Beier	AF	Vachon (1950)
<i>Acanthophorus</i> sp.			

Table 2. Continued.

Carrier	Pseudoscorpion	Region	References
<i>Acrocinus longimanus</i>	<i>Cordyloderes</i> sp.	CA	Muchmore (1971)
<i>Acrocinus longimanus</i>	<i>Cordyloderes costaricensis</i> Beier	NA	Beier (1947)
<i>Acrocinus longimanus</i>	<i>Cordyloderes nigermanus</i> Hoff	CA	Hoff (1944)
<i>Acrocinus longimanus</i>	<i>Cordyloderes panamensis</i> Hoff	CA	Hoff (1944)
<i>Acrocinus longimanus</i>	<i>Cordyloderes scorioides</i> (Linnaeus)	CA	Ellingsen (1913), Zeh & Zeh (1992a)
<i>Acrocinus longimanus</i>	<i>Cordyloderes scorioides</i> (Linnaeus)	CA	Calvert & Calvert (1917)
<i>Acrocinus longimanus</i>	<i>Cordyloderes scorioides</i> (Linnaeus)	SA	Ellingsen (1905)
<i>Acrocinus longimanus</i>	<i>Lustroderes intermedius</i> (Balzan)	SA	Aguiar & Buhrnheim (1992a)
<i>Acrocinus longimanus</i>	<i>Lustroderes</i> sp.	CA	Muchmore (1971)
<i>Acrocinus longimanus</i>	<i>Parachelifer hubbardi</i> (Banks)	NA	Beier (1948)
<i>Acrocinus longimanus</i>	<i>Parachelifer lativittatus</i> (J. C. Chamberlin)	SA	Aguiar & Buhrnheim (1992a)
<i>Acrocinus longimanus</i>	<i>Dendroderes cyrneus</i> (L. Koch)	EU	Beier (1929)
<i>Assem striatum</i>	<i>Lissocheler superbus</i> (With)	AS	Beier (1948)
<i>Batocera celebiana</i>	<i>Dendroderes cyrneus</i> (L. Koch)	EU	Kew (1929)
<i>Callidium variabile</i>	<i>Titanatennus kibweziensis</i> Beier	AF	Beier (1948)
<i>Coelodon servum</i>	<i>Parachelifer persimilis</i> (Banks)	NA	Muchmore (1971)
<i>Ergates spiculatus</i>	<i>Lustroderes grossus</i> (Banks)	NA	Muchmore (1971)
<i>Ergates spiculatus</i>	<i>Cordyloderes scorioides</i> (Linnaeus)	SA	Aguiar & Buhrnheim (1992a)
<i>Macrodontia cervicornis</i>	<i>Titanatennus equester</i> (With)	AF	Beier (1948)
<i>Macrotoma natala</i>	<i>Dendroderes cyrneus</i> L. Koch	EU	Jones (1978)
<i>Phymatodes testaceus</i>	<i>Lechytia chthoniformis</i> (Balzan)	SA	Aguiar & Buhrnheim (1992a)
<i>Stenodontes spinibarbis</i>	<i>Lustroderes intermedius</i> (Balzan)	SA	Aguiar & Buhrnheim (1992a)
<i>Stenodontes spinibarbis</i>	<i>Neochetridium corticum</i> (Balzan)	SA	Aguiar & Buhrnheim (1991)
<i>Tragocephala</i> sp.	<i>Paratennoides pallidus</i> (Balzan)	AF	Beier (1948)
<i>Tragosoma deparius</i>	<i>Parachelifer persimilis</i> (Banks)	NA	Muchmore (1971)
Unknown	<i>Dendroderes instabilis</i> (J. C. Chamberlin)	NA	Chamberlin (1934)
Unknown	<i>Lissocheler</i> sp.	AS	Beier (1948)
Unknown	<i>Lustroderes</i> sp.	NA	Muchmore (1971)
Unknown	<i>Metatennus superior</i> Muchmore	AS	Muchmore (1971)
Unknown	Unknown	AS	Muchmore (1972)
Unknown	Unknown	NA	Anonymous (1986)
Cicindelidae			
<i>Cicindela hirticollis</i>	<i>Parachelifer longipalpus</i> Hoff	NA	Muchmore (1971)
Cleridae			
<i>Thanosinus formicarius</i>	<i>Lanproderes nodosus</i> (Schränk)	EU	Trappen (1906)
Coccinellidae			
<i>Chilocorus</i> sp.	<i>Oratennus brevidigitatus</i> Beier	AF	Beier (1940)
Curculionidae			
<i>Rhynchophorus palmarum</i>	<i>Lustroderes argentinus</i> (Thorell)	SA	Beier (1948)
Elaeteridae			

Table 2. Continued.

Carrier	Pseudoscorpion	Region	References
<i>Alaus oculatus</i>	<i>Americhernes oblongus</i> (Say)	NA	Leidy (1877)
<i>Alaus maculatus</i>	<i>Americhernes oblongus</i> (Say)	NA	Hagen (1879)
<i>Chalcolepidius rugatus</i>	<i>Parachernes nigrimanus</i> Beier	CA	Beier (1948)
<i>Chalcolepidius rugatus</i>	<i>Parachernes setosus</i> Beier	CA	Beier (1948)
<i>Pyrophorus phosphoreus</i>	<i>Lustrochernes argentinus</i> (Thorell)	SA	Beier (1948)
Lampyridae			
<i>Photuris</i> sp.	<i>Paratemnoides elongatus</i> (Banks)	NA	Lloyd <i>et al.</i> (1974), Lloyd & Muchmore (1975)
Passalidae			
<i>Passalus abortivus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus aff. coarctatus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus convexus</i>	<i>Americhernes</i> aff. <i>incertus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus convexus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus convexus</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin & R. V. Chamberlin	SA	Aguar & Buhrnheim (1992b)
<i>Passalus elfriedae</i>	<i>Americhernes</i> aff. <i>incertus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus elfriedae</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus elfriedae</i>	<i>Parawithius gracilimanus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus elfriedae</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin & R. V. Chamberlin	SA	Aguar & Buhrnheim (1992b)
<i>Passalus interruptus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus interruptus</i>	<i>Parawithius gracilimanus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus interruptus</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin & R. V. Chamberlin	SA	Aguar & Buhrnheim (1992b)
<i>Passalus interstitialis</i>	<i>Lustrochernes intermedius</i> (Balzan)	SA	Aguar & Buhrnheim (1992b)
<i>Passalus interstitialis</i>	<i>Parawithius gracilimanus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus interstitialis</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin & R. V. Chamberlin	SA	Aguar & Buhrnheim (1992b)
<i>Passalus latifrons</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus latifrons</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin & R. V. Chamberlin	SA	Aguar & Buhrnheim (1992b)
<i>Passalus punctiger</i>	<i>Lustrochernes argentinus</i> (Thorell)	SA	Beier (1948)
<i>Passalus rhodocanthopoides</i>	<i>Lustrochernes intermedius</i> (Balzan)	SA	Aguar & Buhrnheim (1992b)
<i>Passalus rhodocanthopoides</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)
<i>Passalus rhodocanthopoides</i>	<i>Parawithius gracilimanus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus rhodocanthopoides</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin & R. V. Chamberlin	SA	Aguar & Buhrnheim (1992b)
<i>Passalus unicornis</i>	<i>Americhernes</i> aff. <i>incertus</i> Mahnert	SA	Aguar & Buhrnheim (1992b)
<i>Passalus unicornis</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguar & Buhrnheim (1992b)



Table 2. Continued.

Carrier	Pseudoscorpion	Region	References
<i>Passalus unicornis</i>	<i>Parawithius gracilimanus</i> Mahnert	SA	Aguiar & Buhrnheim (1992b)
<i>Passalus unicornis</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin et R. V. Chamberlin	SA	Aguiar & Buhrnheim (1992b)
<i>Passalus variipylus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguiar & Buhrnheim (1992b)
<i>Venturinus platyrhynus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguiar & Buhrnheim (1992b)
<i>Venturinus transversus</i>	<i>Lustrochernes</i> aff. <i>reimoseri</i> Beier	SA	Aguiar & Buhrnheim (1992b)
<i>Venturinus transversus</i>	<i>Tridenchthonius mexicanus</i> J. C. Chamberlin et R. V. Chamberlin	SA	Aguiar & Buhrnheim (1992b)
Platypodidae			
<i>Platypus</i> sp.	<i>Dolichowithius medifasciatus</i> Mahnert	SA	Aguiar <i>et al.</i> (1992)
Unknown	Unknown	CA	Poinar (unpublished observation)
Prostomidae			
<i>Prostomis mandibularis</i>	<i>Lechytia</i> sp.	NA	Muchmore (1971)
Staphylinidae			
<i>Staphylinus fubomaculatus</i>	<i>Incachernes mexicanus</i> Beier	NA	Beier (1933)
Unknown	<i>Dendrochernes instabilis</i> (J. C. Chamberlin)	CA	Muchmore (1971)
Unknown	<i>Dendrochernes cyrneus</i> (L. Koch)	EU	Lohmander (1939)c
Diptera			
Anthomyiidae			
<i>Anthomyia</i> sp.	<i>Lamprochernes nodosus</i> (Schrank)	EU	Gerstäcker (1859)
<i>Hylemyia floralis</i>	<i>Allochernes peregrinus</i> Lohmander	EU	Lohmander (1939)
<i>Ophyra leucostoma</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Graham-Smith (1916)
<i>Pegomya affinis</i>	<i>Pselaphochernes scorpoides</i> (Hermann)	NA	Muchmore (1971)
<i>Pegomya apicalis</i>	<i>Hesperochernes</i> sp.	NA	Muchmore (1971)
<i>Pegomya apicalis</i>	<i>Lamprochernes savignyi</i> (E. Simon)	AF	Beier (1948)
<i>Pegomya apicalis</i>	<i>Lamprochernes savignyi</i> (E. Simon)	EU	Beier (1948)
Calliphoridae			
<i>Calliphora erythrocephala</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Leydig (1867)
<i>Calliphora vomitoria</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Donovan (1797)
<i>Calliphora vomitoria</i>	<i>Chelifer cancröides</i> (Linnaeus)	EU	Donovan (1797)
<i>Calliphora</i> sp.	<i>Suborder Cheliferinea</i>	SA	Berg (1893)
<i>Lucilia caesar</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Masan & Kristofik (1992)
<i>Lucilia</i> sp.	<i>Lustrochernes argentinus</i> (Thorell)	SA	Beier (1948)
Culicidae			
<i>Aedes sticticus</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Minar (1966)
Drosophilidae			
<i>Drosophila funebris</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Vachon (1940, 1954a)
Heleomyzidae			

Table 2. Continued.

Carrier	Pseudoscorpion	Region	References
<i>Amoebalaria deffesa</i>	<i>Chelodamus</i> sp.	NA	Muchmore (1971)
<i>Amoebalaria</i> sp.	<i>Hesperocheernes taniae</i> Beier	NA	Muchmore (1971)
<i>Tephrochlamis canescens</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Burr (1919)
Lonchaeidae			
<i>Lonchaea chorea</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Jones (1978)
<i>Lonchaea chorea</i>	<i>Lamprocheernes scorpoides</i> (Hermann)	EU	Jones (1978)
<i>Lonchaea latitons</i>	<i>Rhacochelifer similis</i> Beier	AF	Vachon (1940)
<i>Lonchaea vaginalis</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Graham-Smith (1916)
<i>Lonchaea</i> sp.	<i>Lamprocheernes savignyi</i> (E. Simon)	EU	Jones (1978)
Muscidae			
<i>Fannia canalicularis</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Macrae (1869), Zielke (1969)
<i>Hydrotaea similis</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Masan & Kristofik (1992)
<i>Musca autumnalis</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Jones (1970, 1978)
<i>Musca corvina</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Graham-Smith (1916)
<i>Musca domestica</i>	<i>Chelifer cancrroides</i> (Linnaeus)	EU	Nonidez (1917)
<i>Musca domestica</i>	<i>Chernes sanbornii</i> Hagen	NA	Webster (1897)
<i>Musca domestica</i>	<i>Dinocheirus panzeri</i> (C. L. Koch)	EU	Preudhomme (1873)
<i>Musca domestica</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Poda (1761), Carl (1994)
<i>Musca domestica</i>	<i>Neobisium sylvaticum</i> (C. L. Koch)	EU	PRESENT STUDY
<i>Musca domestica</i>	<i>Pselaphocheernes scorpoides</i> (Hermann)	EU	Jones (1978)
<i>Musca domestica</i>	<i>Lamprocheernes</i> sp.	NA	Muchmore (1971)
<i>Musca domestica</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Beier (1948)
<i>Musca meteorica</i>	<i>Chelifer cancrroides</i> (Linnaeus)	EU	Beier (1948)
<i>Musca vicina</i>	<i>Muscitcheernes katoi</i> (Morikawa)	AS	Morikawa (1960)
<i>Musca</i> ?	<i>Pselaphocheernes</i> sp.	NA	Muchmore (1971)
<i>Stomoxys calcitrans</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Lukis (1831), Schiner (1872)
<i>Stomoxys calcitrans</i>	<i>Lamprocheernes savignyi</i> (E. Simon)	EU	Jones (1978)
<i>Stomoxys</i> sp.	<i>Pselaphocheernes scorpoides</i> (Hermann)	EU	Graham-Smith (1916)
Unknown	<i>Dinocheirus sicarius</i> J. C. Chamberlin	NA	Chamberlin (1952)
Neriidae			
<i>Odomolozus longicornis</i>	<i>Dinocheirus arizonensis</i> (Banks)	NA	Zeh & Zeh (1992c)
Otitidae			
<i>Physiphora demandata</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Clements (1987)
<i>Ulidia erythrophthalma</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Low (1866)
<i>Ulidia erythrophthalma</i>	<i>Allocheernes wideri</i> (C. L. Koch)	EU	Beier (1948)
Pantophthalmidae			
<i>Pantophthalmus tabaninus</i>	<i>Semeiocheernes armiger</i> (Balzan)	CA	Zeh & Zeh (1992b)
Scenopinidae			
<i>Omphrale fenestralis</i>	<i>Lamprocheernes nodosus</i> (Schrank)	EU	Vachon (1940)

Table 2. Continued.

Carrier	Pseudoscorpion	Region	References
Sphaeroceridae			
<i>Lepiocera sylvatica</i>	<i>Pselaphoernes scorioides</i> (Hermann)	EU	Cuthbertson (1982)
Unknown	<i>Pselaphoernes scorioides</i> (Hermann)	EU	Weygoldt (1969)
Stratiomyidae			
<i>Hoplitès bipinosus</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Vachon (1940)
Syrphidae			
<i>Brachypalpus laphriformis</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Gerstäcker (1859)
<i>Eristalis arbustorum</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Jones (1978)
<i>Sargus iridatus</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Jones (1978)
<i>Volucella zonaria</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Berland (1932)
Tabanidae			
<i>Tabanus</i> sp.	Suborder Cheliferinea	SA	Berg (1893)
Unknown (Dexinae)	<i>Hesperochernes pallipes</i> (Banks)	NA	Banks (1895)
Unknown	<i>Parachernes argentatopunctatus</i> (Ellingsen)	SA	Beier (1948)
Unknown	<i>Lamprochernes savignyi</i> (E. Simon)	AU	Ellingsen (1913)
Tachinidae			
<i>Tachina larvarum</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Beier (1948)
<i>Tachina larvarum</i>	<i>Chelifer cancrroides</i> (Linnaeus)	EU	Beier (1948)
Tipulidae			
<i>Ctenophora pectinicornis</i>	<i>Anthrenochernes stellae</i> Lohmander	EU	Gardenfors & Wilander (1995)
<i>Ctenophora pectinicornis</i>	<i>Lamprochernes nodosus</i> (Schrank)	EU	Wagner (1892)
<i>Ctenophora pectinicornis</i>	<i>Chernes cimicoides</i> (Fabricius)	EU	Wagner (1892)
<i>Limonia decemmaculata</i>	<i>Pselaphochernes scorioides</i> (Hermann)	EU	Jones (1978)
<i>Tipula uncinata</i>	<i>Syarinus obscurus</i> (Banks)	NA	Muchmore (1971)
<i>Tipula</i> sp.	<i>Syarinus obscurus</i> (Banks)	NA	Beier (1948)
<i>Zelandotipula</i> sp.	<i>Chernetidae</i>	SA	Mathiesen & Hahn (1981)
Unknown	" <i>Lamprochernes loewi</i> Hagen" (nomen nudum)	CA	Hagen (1879)
Unknown	<i>Lamprochernes savignyi</i> (Simon)	NZ	Muchmore (1971)
Unknown	<i>Lamprochernes nodosus</i> (Schrank)	AS	Redikozev (1936)
Unknown	<i>Semeiochernes militaris</i> Beier	SA	Mahnert (1987)
Unknown	"Pseudoscorpionida"	PA (on boat)	Guilmette <i>et al.</i> (1970)
LEPIDOPTERA			
Arctiidae			
<i>Spilosoma ratrayi</i>	<i>Stenowithius bayoni</i> (Ellingsen)	AF	Beier (1932)
Cocytidae			
<i>Cocytius duponchel</i>	<i>Neocheiridium triangulare</i> Mahnert et Aguiar	SA	Mahnert & Aguiar (1986)
Geometridae			
<i>Sterrhia aversata</i>	<i>Rhacochelifer maculatus</i> (L. Koch)	EU	Vachon (1953)

Table 2. Continued.

Carrier	Pseudoscorpion	Region	References
Hesperidae			
<i>Pelopides tridens</i>	<i>Macrochelifer bornoensis</i> (Ellingsen)	AS	Beier (1948)
Lymantriidae			
<i>Lymantria monacha</i>	<i>Chelifer cancroides</i> (Linnaeus)	EU	Ketterer (1955)
<i>Spilosoma ratsayi</i>	<i>Stenowithius bayoni</i> (Ellingsen)	AF	Beier (1932)
Noctuidae			
<i>Acronycta grisea</i>	<i>Apocheiridium</i> sp.	NA	Treat (1956)
<i>Acronycta morula</i>	<i>Apocheiridium</i> sp.	NA	Muchmore (1971)
<i>Acronycta ovata</i>	<i>Apocheiridium</i> sp.	NA	Treat (1956)
<i>Caenurgina crassiuscula</i>	<i>Dinocheirus pallidus</i> (Banks)	NA	Fenstermacher (1959)
<i>Catocala neogama</i>	<i>Apocheiridium</i> sp.	NA	Muchmore (1971)
<i>Rhyacia augur</i>	<i>Lamprochernes chyzeri</i> (Tömösvary)	EU	Vachon (1954)
Riodinidae			
<i>Eurybia</i> sp.	Unknown	CA	De Vries (personal communication)
<i>Synargus</i> sp.	Unknown	CA	De Vries (personal communication)
Unknown	<i>Withius piger</i> (Simon)	AF	Beier (1930)
Microlepidoptera	Unknown	EU	Berland (1932)
Hymenoptera			
Apidae			
<i>Apis mellifera</i>	<i>Chelifer cancroides</i> (Linnaeus)	NA	Filleul (1922)
<i>Apis mellifera</i>	<i>Ellingseniulus fulleri</i> (Hewitt et Godfrey)	AF	Hewitt & Godfrey (1929)
<i>Apis mellifera</i>	<i>Ellingseniulus hendrickxi</i> Vachon	AF	Vachon (1954)
<i>Apis mellifera</i>	<i>Ellingseniulus sculpturatus</i> (Lewis)	AF	Ward (1887)
<i>Apis mellifera</i>	<i>Paratennoides nidificator</i> (Balzan)	CA	Muchmore (1971)
Unknown	<i>Cordylochernes scorioides</i> (Linnaeus)	SA	Tömösvary (1884)
"Wild bee"	<i>Lamprochernes</i> sp.	NA	Muchmore (1971)
Aulacidae			
<i>Pristaulacus niger</i>	<i>Dendrochernes</i> cf. <i>morosus</i> (Banks)	NA	Haack & Wilkinson (1986)
Formicidae			
<i>Acromyrmex lundii</i>	<i>Sphaenochernes bruchi</i> (Mello-Leitão)	SA	Mello-Leitão (1927)
<i>Oecophylla smaragdina</i>	<i>Cheliferinea</i> sp.	AS	Green (1908)
Ichneumonidae			
<i>Ephialtes mesocentrus</i>	<i>Chernes cimicoides</i> (Fabricius)	EU	Jones (1978)
Siricidae			
<i>Tremex columba</i>	<i>Dendrochernes</i> sp.	NA	Muchmore (1971)

flight, carrying the pseudoscorpion along with it. Jones (1978) concurred and felt that phoretic behavior in pseudoscorpions was a modification of their original hunting instinct, which caused them to seize large animals that passed by.

We agree that phoresy in pseudoscorpions can be motivated by hunger, however not hunger for the carrier, but rather a desire to reach a new habitat that has a greater nutritional potential than the present location. Thus, in the great majority of, if not in all, cases involving arthropod phoresy in pseudoscorpions, the advantage and ultimate goal of the carrier is to reach a new habitat (i.e., dispersal). This is not to say that under other circumstances, the pseudoscorpion will not turn around and attack its potential carrier. An interesting case has been documented where a neriid fly, *Odontolorozus longicornis*, serves as both carrier and prey to *Dinocheirus arizonicus* (Zeh & Zeh 1992c). In this situation, the pseudoscorpion exhibited a clear distinction between phoretic and predatory behavior towards the fly. When initiating a phoretic relationship, the arachnid would grasp the fly's hind trochanter with its chela; when exhibiting predatory behavior, however, it would secure the fly with its pedipalpal chela in order to inject the victim with venom.

A difference between predatory and dispersal responses was demonstrated by Cuthbertson (1984) with a captured *Dendrochernes cyrneus*. When a female pseudoscorpion grasped a small piece of paper, as it would an arthropod appendage, it entered a quiescent state, which the author referred to as catalepsy. While in this state, any predatory activity was out of the question, showing that this type of grasping reflex was adapted for dispersal only. Additional observations on sex selection of carriers with *Semeiochernes armiger* also support the hypothesis that phoretic behavior is for dispersal only (Zeh & Zeh 1992b). The pseudoscorpions would only persist in grasping female timber flies (*Pantophthalmus tabaninus*), but quickly released their hold on enclosing males. This was considered an adaptation for dispersal, since only the female flies visit trees suitable for pseudoscorpion establishment. If phoresis in this association was initiated as a predatory response, then both sexes of the fly would be equally treated.

These reports demonstrate that the response of pseudoscorpions resulting in phoresis is quite separate from predatory actions and that phoretic behavior does not have predation as its basis.

That phoretic behavior between arthropods and pseudoscorpions can actually disperse the rider over long distances, even trans-oceanic, is suggested by the presence of *Ediogaryops pumilis* on Little Cayman Island in the Caribbean (Harvey 1985) and at least 11 species of 7 families on Krakatoa (Harvey 1986). In addition, pseudoscorpions have been found among airborne insect specimens collected over water on ships (Guilmette *et al.* 1970).

In both the extinct and extant examples of phoresy with platypodid beetles (see Tables 1 & 2), the pseudoscorpion probably fed upon mites, insect eggs, young larvae, and possibly nematodes that occurred in the tunnels of the beetle (mites are also phoretic on platypodid beetles, but there are no reported examples of double phoresis involving mites and pseudoscorpions as in the harlequin beetle). That these phoretic associations may be highly specialized is shown by the constant presence of a specific pseudoscorpion on a particular host, undoubtedly related to the fact that both the carrier and pseudoscorpion occur in the same developmental habitat (soil, under bark, beetle galleries, rotting debris, etc.). This type of co-evolution probably developed over a period of time and was related to the survival strategy of the pseudoscorpion, which seems to gain most from

the association, since adaptations to survival in transitory habitats must be accompanied by effective dispersal mechanisms. Indirect benefits for the carrier could occur if it or its young fed on the developmental stages of the arachnid or if the pseudoscorpion fed on enemies of various stages of the carrier.

It is clear from Table 2 that most reported extant phoretic associations between pseudoscorpions and arthropods involve carriers in the Diptera. And most of these carriers have breeding stages in the soil, rotting debris, or wood galleries, habitats that would be attractive to pseudoscorpions. The next largest order of carriers belongs to representatives of the Coleoptera, with many of these beetles living in habitats similar to the fly carriers.

A concept in evolutionary biology which has been substantiated by a wide range of data is that of behavioral fixity (Boucot 1990). This maxim states that once a successful type of behavior has become established in an organism, it continues until the organism becomes extinct. It also implies that the behavior of extinct organisms will be similar to that of extant representatives in the same genus and often family. Representatives of all four families of pseudoscorpions mentioned in the present study as exhibiting phoresy in the fossil record have similar associations today with same groups of carriers. Even the method of attachment is similar in modern representatives to those of the fossils. For instance, in both the fossil and extant examples of pseudoscorpions carried by platypodid beetles, the pseudoscorpion holds on to an area under the posterior edge of the elytra with a single chela (compare Fig. 2 in the present study with fig. 1 of Aguiar *et al.* 1992). Thus, "modern" phoresy in pseudoscorpions has been in effect for at least 40 million years and probably much longer. Only one of the families of carriers in fossil associations (Braconidae) is not represented in reported extant associations. However, it is premature to say that the association is absent today: it probably has simply not yet been reported. The length of time that these associations have existed suggests that phoretic behavior may indeed be obligatory for the survival of certain species of pseudoscorpions, thus resulting in sophisticated capabilities of detection, attachment, and release regarding the carrier.

In the fossil example of platypodid carrier in Dominican amber (Fig. 2), the beetle is carrying phoretic nematodes as well as a pseudoscorpion. Microbivorous and phytophagous nematodes are often carried by arthropods in phoretic relationships, with the same advantages to the nematode as is obtained for phoretic pseudoscorpions, namely transportation to a new environmental with nourishment and breeding potential (Poinar 1983).

### Acknowledgements

The authors thank Roberta Poinar for comments on the present version. We also thank P. J. De Vries for unpublished records of pseudoscorpion phoresy on riordinid butterflies in Central America. One of us (BPMC) is grateful for the financial help, received from the Serbian Ministry of Science and Technology (Grant 03E03).

### References

- Aguiar, N. O. & P. F. Buhrnheim, 1991. Pseudoscorpiones foreticos de *Stenodontes spinibarbis* (Lin., 1758) (Coleoptera) e redes cricao *Lechytiia chthoniiformis* (Balzan, 1890) (Pseudoscorpiones,

- Chthoniidae) da Ilha de Maraca-Roraima. *Acta Amazonica*, **21**: 425-433.
- Aguiar, N. O. & P. F. Buhrnheim, 1992a. Pseudoscorpiones foreticos de Cerambycidae (Coleoptera) e ocorrência de *Parachelifer* Chamberlin, 1932 (Pseudoscorpiones, Cheliferidae) no Amazonia. *Bol. Mus. para Emilio Gouldi, Ser. zool.*, **812**: 343-348.
- Aguiar, N. O. & P. F. Buhrnheim, 1992b. Pseudoscorpiones (Arachnida) em associacao foretica com Passalidae (Insectos, Coleoptera) no Amazonas, Brasil. *Amazoniana*, **12**: 187-205.
- Aguiar, N. O., J. V. Silva & P. F. buhrnheir, 1992. *Dolichowithius mediofasciatus* Mahnert, 1979 (Arachnida, Pseudoscorpiones, Withiidae) em foresia com Platypodidae (Insecta, Coleoptera) no Amazonas, Brasil. *Amazoniana*, **12**: 181-185.
- Anonymous, 1986. Buggy ride. *Nat. Hist.*, **3**: 100-101.
- Banks, N., 1895. Chernetid attached to a fly. *Ent. News*, **6**: 115.
- Beier, M., 1929. Alcuni Pseudoscorpioni raccolti da C. Menozzi. *Boll. Soc. ent. Ital.*, **61**: 154-156.
- Beier, M., 1930. Die Pseudoscorpione des Wiener naturhistorischen Museums. *Ann. naturhistor. Mus. Wien*, **44**: 199-222.
- Beier, M., 1932. Zur Kenntnis der Cheliferidae (Pseudoscorpionidea). *Zool. Anz.*, **100**: 53-67.
- Beier, M., 1933. Pseudoscorpione aus Mexico. *Zool. Anz.*, **104**: 91-101.
- Beier, M., 1937. Pseudoscorpione aus dem baltischen Bernstein. Festschrift fur Prof. Dr. Embrik Strand (Riga, Latvia), **2**: 302-316.
- Beier, M., 1940. Die Pseudoscorpionidenfauna der landfernen Inseln. *Zool. Jahrb. Syst.*, **74**: 161-192.
- Beier, M., 1947. Pseudoskorpione im baltischen Bernstein usw. *Mikroskopie*, **1**: 188-199.
- Beier, M., 1948. Phoresie und Phagophilie bei Pseudoscorpionen. *Oesterr. zool. Zeitschr.*, **1**: 441-497.
- Beier, M., 1963. Ordnung Pseudoscorpionidea (Afterskorpione). In: *Bestimmungsbucher zur Bodenfauna Europas*, **1**: 1-313, Berlin.
- Berg, C., 1893. Pseudoscorpionidenkniffe. *Zool. Anz.*, **16**: 446-448.
- Berland, L., 1932. Les Arachnides. Mus. Nat. Hist. Nat., Paris.
- Boucot, A. J., 1990. Evolutionary Paleobiology of Behavior and Coevolution. Elsevier, Amsterdam.
- Burr, A., 1919. Epizoisme des Pseudoscorpions. *Assoc. Phil. Alsace-Lorraine*, **4**: 24-26.
- Calvert, A. S. & P. P. Calvert, 1917. A year of Costa Rican Natural History. MacMillan Co., New York.
- Carl, M., 1994. Injurious effects on the exoskeleton of *Musca domestica* L. (Diptera) of phoresy by *Lamprochernes nodosus* (Schrank) (Pseudoscorpiones, Chernetidae) and the possible functional significance of accessory teeth on the chelal fingers. *Bull. Br. arachnol. Soc.*, **9**: 246-248.
- Chamberlin, J. C., 1934. On two species of false scorpions collected by birds in Montana, with notes on the genus *Dendrochernes*. *Pan-Pacific Entomol.*, **10**: 125-132.
- Charberlin, J. C., 1952. New and little-known false scorpions (Arachnida, Chelonethida) from Monterey County, California. *Bull. Amer. Mus. Nat. Hist.*, **99**: 259-312.
- Clements, D. K., 1987. A case of phoresy in the pseudoscorpion *Lamprochernes nodosus* (Schrank) (Pseudoscorpiones: Chernetidae). *Ent. Month. Mag.*, **123**: 222.
- Cloudsley-Thompson, J. L., 1956. Notes on Arachnida. 25. An unusual case of phoresy by false scorpions. *Ent. Month. Mag.*, **92**: 71.
- Cuthbertson, D. R., 1982. Pseudoscorpions attached to flies: phoresy or predation? *Newsl. Br. Arachnol. Soc.*, **34**: 4.
- Cuthbertson, D. R., 1984. Catalepsy and phoresy in pseudoscorpions. *Newsl. Br. Arachnol. Soc.*, **39**: 9.
- Fenstermacher, J. D., 1959. Survey of the pseudoscorpions of Michigan. M. S. Thesis, Michigan State University, 1-43.
- Donovan, E., 1797. The Natural History of British Insects. 6. Donovan, London.
- Dunkle, S. W., 1984. First record of pseudoscorpions phoretic on dragonflies. *Not. Odonatol.*, **2**: 48.
- Durden, L. A., 1991. Pseudoscorpions associated with mammals in Papua New Guinea. *Biotropica*, **23**: 204-206.
- Ellingsen, E., 1905. Pseudoscorpions from South America collected by Dr. A. Borelli, A. Bertoni De Winkelried, and Prof. Goeldi. *Boll. Mus. Zool. Anat. Comp. Univ. Torino*, **500**: 1-17.
- Ellingsen, E., 1913. Note on some Pseudoscorpions in the British Museum. *Ann. Mag. Nat. Hist.*, **11**:

- 451-455.
- Filleul, E. E., 1922. The false scorpion. *Bee World*, **4**: 140.
- Gardenfors, U. & P. Wilander, 1955. Ecology and phoretic habits of *Anthrenochernes stellae* (Pseudoscorpionida, Chernetidae). *Bull. Br. arachnol. Soc.*, **10**: 28-30.
- Gerstacker, A. 1859. Berichte uber die wissenschaftlichen Leistungen im Gebiete der Entomologie. Berlin.
- Graham-Smith, G., 1916. Observations of the habits and parasites of common flies. *Parasitology*, **4**: 514-523.
- Green, E. E., 1908. Curious habits of Chelifers. *Zoologist*, **12**: 159-160.
- Guilmette, J. E., Jr. E. P. Holzapfel & D. M. Tsuda, 1970. Trapping of air-borne insects on ships in the Pacific. Part 8. *Pacific Insects*, **12**: 303-325.
- Haack, R. A. & R. C. Wilkinson, 1986. Phoresy by *Dendrochernes* pseudoscorpions on Cerambycidae (Coleoptera) and Aulacidae (Hymenoptera) in Florida. *Amer. Midland Nat.*, **117**: 369-373.
- Hagen, H., 1879. Hohlen-Chelifer in Nord America. *Zool. Anz.*, **2**: 399-400.
- Harvey, M. S., 1985. The systematics of the family Sternophoridae (Pseudoscorpionida). *J. Arachnol.*, **13**: 141-209.
- Harvey, M. S., 1986. The systematics and biology of Pseudoscorpions. Austral. Arachnol., Austral. Entom. Soc., Brisbane, 75-85.
- Hewitt, J. & R. Godfrey, 1929. South African pseudoscorpions of the genus *Chelifer* Geoffroy. *Ann. Natal Mus.*, **6**: 305-336.
- Hoff, C. C., 1944. New pseudoscorpions of the subfamily Lamprochernetinae. *Amer. Mus. Nov.*, **1271**: 1-12.
- Hoff, C. C., 1947. Two new pseudoscorpions of the subfamily Lamprochernetinae from Venezuela. *Zoologica, Sc. Contr. N. Y. Zool. Soc.*, **32**: 61-64.
- Hoff, C. C., & D. T. Jennings, 1974. Pseudoscorpions phoretic on a spider. *Entomol. News*, **85**: 21-22.
- Jones, P. E., 1970. *Lamprochernes nodosus* (Schrank)-an example of phoresy in pseudoscorpions. *Bull. Br. Arach. Soc.*, **1**: 118-119.
- Jones, P. E., 1978. Phoresy and commensalism in British Pseudoscorpions. *Proc. Brit. Ent. nat. Soc.*, 1978: 90-96.
- Joseph, G., 1882. Systematisches Verzeichnis der in den Tropfsteingrotten von Krain einheimischen Arthropoden. *Berl. ent. Zeitschr.*, **26**: 1-50.
- Kaestner, A., 1970. Invertebrate Zoology. Arthropod relatives, Chelicerata, Myriopoda. Vol. 11. John Wiley & Sons, New York.
- Ketterer, C.-EM., 1955. A propos d'un cas de phoresie chez *Chelifer cancroides* (Pseudoscorpionidea). *Bull. Murith.*, **72**: 93-97.
- Kew, H. W., 1929. Notes on some Coleoptera and a *Chelifer*, observed on a Richmond Park Oak after nightfall. *Ent. Month. Mag.*, **15**: 83-86.
- Leidy, J., 1877. Remarks on the seventeen-year locust, the Hessian fly and an *Chelifer*. *Proc. Acad. Nat. Sci. Philadelphia*, 1877: 260-261.
- Legg, G., 1975. The possible significance of spermathecae in pseudoscorpions (Arachnida). *Bull. Br. Arach. Soc.*, **3**: 91-95.
- Leydig, F., 1867. Skizze zu einer Fauna Tubingensis. Stuttgart.
- Lloyd, J. E., S. Correale & W. B. Muchmore, 1974. Pseudoscorpions phoretic on fireflies. *Fla. Entomol.*, **57**: 381.
- Lloyd, J. E. & W. B. Muchmore, 1975. Pseudoscorpions phoretic on fireflies. 11. *Fla. Entomol.*, **58**: 241-242.
- Lohmander, H., 1939. Zwei neue Chernetiden der nordwesteuropischen Fauna. *Goteborgs Vet. Vitt. Samh. Handl.*, **B 6/11**: 1-11.
- Low, F., 1866. Zoologische Notizen. 1. Arachnoidea. *Verh. zool.-bot. Ges. Wien*, **16**: 944-945.
- Lukis, F. C., 1831. A lobster-like insect attacking the leg of a house-fly. *Ann. Mag. nat. Hist.*, **4**: 283-284.



- Macrae, G., 1869. Notice. *Hardwicks Sc. Gossip*, 1: 223.
- Mahnert, V., 1987. Neue oder wenig bekannte, vorwiegend mit Insekten vergesellschaftete Pseudoscorpione (Arachnida) aus Südamerika. *Mitt. Schweiz. Ent. Ges.*, **60**: 403-416.
- Mahnert, V. & N. O. Aguiar, 1986. Wieder Beschreibung von *Neoecheiridium corticum* (Balzan, 1890) und Beschreibung von zwei neuen Arten der Gattung aus Südamerika (Pseudoscorpiones, Cheiridiidae). *Mitt. Schweiz. Ent. Ges.*, **59**: 499-509.
- Masan, P. & J. Kristofik, 1992. Phoresy of some arachnids (Acarina and Pseudoscorpionea) on synanthropic flies (Diptera) in the south Slovakia. *Biologia (Bratislava)*, **47**: 87-96.
- Matthiesen, F. A. & N. S. Hahn, 1981. Foresia em pseudo-escorpioes brasileiros. *Ciencia e Cultura*, **33**: 689-690.
- Menge, A., 1855. Über die Scheerenspinnen, Chernetidae. *Schr. naturf. Ges. Danzig*, **5**: 1-41.
- Mello-Leitão, M., 1927. Dois interessantes arachnidos myrmecophilos. *Physis Buenos-Aires*, **8**: 228-237.
- Minar, J., 1966. Phoresia of *Damalinia* (C.) *meyeri* (Mallophaga) and *Lamprochernes nodosus* (Pseudoscorpionidea) on mosquitoes *Aedes sticticus* (Culicidae). *Folia Parasitologica*, **13**: 270-273.
- Morikawa, K., 1960. Systematic studies of Japanese pseudoscorpions. *Mem. Echime Univ., Sect. 2 (Sci.), Ser. B (Biol.)*, **4**: 85-172.
- Muchmore, W. B., 1971. Phoresy by North and Central American Pseudoscorpions. *Proc. Rochester Acad. Sci.*, **12**: 79-97.
- Muchmore, W. B., 1972. A phoretic *Metatemnus* (Pseudoscorpionida, Atemnidae) from Malaysia. *Entomol. News*, **83**: 11-14.
- Nonidez, J. F., 1917. Pseudoscorpions de España. *Trab. Mus. Nac. Cienc. Nat. Madrid*, **32**: 1-46.
- Poda, N., 1761. *Insecta Musei Graecensis, Quae in Ordines, Genera et Species Iuxta Systema Naturae Caroli Linnaei*. Athens.
- Poinar, JR. G. O., 1983. *Natural History of Nematodes*. Prentice Hall, Englewood Cliffs, New Jersey.
- Poinar, JR. G. O., 1992. *Life in amber*. Stanford University Press, Stanford, CA.
- Preudhomme, D. B., 1873. Note biologique relative aux mœurs des Arachnides du genre *Chelifer*. *Verh. zool. bot. Ges. Wien*, **23**: 36.
- Redikorzev, V. V., 1936. Pseudoscorpiones. In: *Animals of the world-USSR*. Redikorzev, V. V. (Ed.), Academy of Sciences of the USSR, Moscow.
- Ross, A. J., 1997. Insects in amber. *Geology Today*, **13**: 24-28.
- Sankey, J. H. P., 1949. Observations on the food, enemies and parasites of British harvest-spiders (Arachnida, Opiliones). *Entom. Mon. Mag.*, **85**: 246-247.
- Schawaller, W., 1981. Pseudoscorpione (Cheliferidae) phoretisch auf Kanfern (Platypodidae) in Dominikanischen Bernstein (Stuttgarter Bernsteinsammlung: Pseudoscorpionidea und Coleoptera). *Stuttgarter Beitr. Naturk.*, **71B**: 17 pp.
- Schlee, D., 1980. Bernstein-Raritäten., Staatliches Museum für Naturkunde, Stuttgart.
- Schlee, D. & W. Glockner, 1978. Bernstein. *Stuttgarter Beitr. Naturk.*, **8C**: 1-72.
- Shear, W. A., W. Schawaller & P. M. Bonamo, 1989. Record of Paleozoic pseudoscorpions. *Nature*, **341**: 527-529.
- Schiner, T., 1872. Miscellen. Vorkommen von *Chelifer* an Fliegen. *Verh. zool.-bot. Ges. Wien*, **22**: 75-76.
- Spicer, W. W., 1867. Helps to distribution. *Hardwicks Sc. Gossip*, **1**: 244-245.
- Tömösvary, O., 1884. Adatok az Alskorpiók ismeretéhez. *Termesz. Füzetek*, **8**: 16-27.
- Trappen, V., 1906. Sonderbare Jäger. *Soc. Entomol.*, **21**: 52.
- Treat, A. E., 1956. A pseudoscorpion on moths. *Lepidopterists News*, **10**: 87-89.
- Vachon, M., 1940. Remarques sur la phorésie des Pseudoscorpions. *Ann. Soc. Ent. France*, **109**: 1-18.
- Vachon, M., 1947. Nouvelles remarques à propos de la phoresie des Pseudoscorpions. *Bull. Mus. nat. Hist. nat. Paris*, **19**: 84-87.
- Vachon, M., 1950. A propos d'une "association" phorétique: Coléoptère-Acariens-Pseudoscorpions. *Bull. Mus. nat. Hist. nat. Paris*, **22**: 728-733.
- Vachon, M., 1953. Nouveaux cas de phorésie chez les Pseudoscorpions. *Bull. Mus. nat. Hist. nat.*

*Paris*, **25**: 572-575.

- Vachon, M., 1954. Remarques sur un Pseudoscorpion vivant dans les ruches d'Abeilles au Congo Belge. *Ann. Mus. Congo Tervuren, Zool.*, **1**: 284-287.
- Vachon, M., 1954a. Nouvelles captures de Pseudoscorpions (Arachnides) transportés par des insectes. *Bull. Mus. nat. Hist. nat. Paris*, **26**: 590-592.
- Wagner, F. V., 1892. Biologische Notizen. *Zool. Anz.*, **15**: 434-436.
- Ward, J. R., 1887. Bees in Natal: their companions and parasites. *Brit. Bee Journ.*, **15**: 563.
- Webster, F. M., 1897. Pseudoscorpions attached to flies. *Ent. News Philad.*, **8**: 59.
- Weygoldt, P., 1969. The biology of pseudoscorpions. Cambridge, Mass, Harvard University Press.
- Wu, R. J. C., 1996. Secret of a lost world. Dominican amber and its inclusions. Self Published. Santa Domingo, Dominican Republic.
- Zeh, D. W. & J. A. Zeh, 1991. Novel use of silk by the harlequin beetle-riding pseudoscorpion, *Cordylorchernes scorioides* (Pseudoscorpionida: Chernetidae). *J. Arachn.*, **19**: 153-164.
- Zeh, D. W. & J. A. Zeh, 1992a. Dispersal-generated sexual selection in a beetle-riding pseudoscorpion. *Behav. Ecol. Sociobiol.*, **30**: 135-142.
- Zeh, D. W. & J. A. Zeh, 1992b. Emergence of a giant fly triggers phoretic dispersal in the neotropical pseudoscorpion, *Semeiochernes armiger* (Balzan) (Pseudoscorpionida: Chernetidae). *Bull. Br. Arachnol. Soc.*, **9**: 43-46.
- Zeh, D. W. & J. A. Zeh, 1992c. Failed predation or transportation? Causes and consequences of phoretic behavior in the pseudoscorpion *Dinochernes arizonensis* (Pseudoscorpionida: Chernetidae). *J. Insect Behav.*, **5**: 37-49.
- Zeh, D. W. & J. A. Zeh, 1992d. On the function of harlequin beetle-riding in the pseudoscorpion *Cordylorchernes scorioides* (Pseudoscorpionida: Chernetidae). *J. Arachnol.*, **20**: 47-51.
- Zielke, E., 1969. Beobachtungen zur Phoresie bei *Lamprochernes nodosus* (Schränk). *Mitt. Hamburg. Zool. Staatsinst. Zool. Mus. Entomol.*, **3**: 1-3.